Vigoro Ultra Turf™

Weed & Feed

28-3-3

Herbicida y Fertilizante

TREATS 15,000 SQ FT

TRATA 15,000 PIES²

FPO UPC
0 13013 22344 1

Vigoro®

Ultra Turf™

Weed & Feed

28-3-3

Herbicida y Fertilizante

Kills Over 200 Weed Types!

¡Elimina más de 200 Tipos de Malezas!

- ***Controls Dandelion, White Clover, Plantain, Chickweed, Spurge*** *& Many Other Listed Weeds*
  Controla el Diente de León, el Trébol Blanco, el Llantén, la Pamplina, la Lechetrezna *y muchas otras Malezas Enumeradas*
- ***Greens in 72 Hours***
  Proporciona Color Verde en 72 Horas
- ***Contains Composite Timed-Release (CTR®)*****
  Nitrogen for Extended Feeding
  Contiene Fertilizante de Liberación Programada Compuesta (CTR®) para una Nutrición de Mayor Duración**

ACTIVE INGREDIENTS:	
2,4-D	0.64%
Mecoprop-p	0.16%
Dicamba	0.03%
OTHER INGREDIENTS:	99.17%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
12.80 lbs. 2, 4-dichlorophenoxyacetic acid equivalent per ton.
3.20 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
0.60 lbs. dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.
Isomer Specific by AOAC Methods
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

CAS Registry Numbers:
2,4-D CAS# 94-75-7
Mecoprop-p CAS# 16484-77-8
Dicamba CAS# 1918-00-9

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements and First Aid.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN Consulte el panel posterior para obtener Declaraciones de Precaución y Primeros Auxilios adicionales.

Net Wt 48 lb (21.7 kg)

TREATS 15,000 SQ FT Trata 15,000 pies²	APPLY ANYTIME WEEDS ARE ACTIVELY GROWING Aplique en cualquier momento en que las malezas estén creciendo activamente

Vigoro® Ultra Turf™

Weed & Feed
28-3-3
Herbicida y Fertilizante

English

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PROJECT GUIDE

For a thicker, greener, more beautiful lawn, follow ***Vigoro's Annual Lawn Care Program***. It's easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!

SPRING ▸ ▸ ▸ ▸ ▸ FALL ▸ ▸ ▸

Early Spring	*Late Spring*	*Summer*	*Early Fall*	*Late Fall*
CRABGRASS PREVENTER	WEED & FEED	LAWN INSECT CONTROL	LAWN FERTILIZER	WINTERIZER

Vigoro offers complete annual lawn care programs for all areas. See your local store associate or visit ***Vigoro.com*** for more information about a customized program that is right for your area and turf type.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

Vigoro® Ultra Turf™ Weed & Feed 28-3-3 controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

For Use On:

Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, Fescue, Kentucky bluegrass, Perennial ryegrass, St. Augustinegrass, Zoysiagrass

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass:

Apply 3.2 lb of product per 1,000 sq ft of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 lb of product per 1,000 sq ft should be used 30 days after the first application. **Do not apply this product to 'Floratam' St. Augustinegrass in Florida.**

The maximum application rate to turf is 1.1 lb of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

A lawn typically utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.89 lb of nitrogen per 1,000 sq ft. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application. Do not apply if rain is expected within 2 days.

Weeds Controlled

Annual yellow sweetclover, Aster, Austrian fieldcress, Bedstraw, Beggarticks, Betony, Florida, Bindweed, field, Bird vetch, Bitter wintercress, Bittercress, hairy, Bitterweed, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Bloodflower milkweed, Blue lettuce, Blue vervain, Bracted plantain, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull thistle, Bullnettle, Burclover, Burdock, Burning nettle, Burweed, Buttercup, Buttonweed, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catnip, Catsear, Chickweed, common, Chickweed, mouseear, Chicory, Cinquefoil, Clover, crimson, Clover, hop, Clover, red, Clover, strawberry, Clover, sweet, Clover, white, Cockle, Cocklebur, Common mullein, Creeping jenny, Cudweed, Curly dock, Daisy, English, Daisy fleabane, Daisy, oxeye, Dandelion, Dichondra, Dogfennel, Dollarweed, False dandelion, Falseflax, False sunflower, Fiddleneck, Florida pusley, Frenchweed, Galinsoga, Goldenrod, Ground ivy, Gumweed, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Heath aster, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk vetch, Morningglory, Mouseear hawkweed, Mugwort, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oriental cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prairie sunflower, Prickly lettuce, Prickly sida, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purslane, common, Ragweed, Red sorrel, Redroot pigweed, Redstem filaree, Rough cinquefoil, Rough fleabane, Russian pigweed, Russian thistle, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smooth dock, Smooth pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny amaranth, Spiny cocklebur, Spiny sowthistle, Spotted catsear, Spotted spurge, Spurweed, Stinging nettle, Strawberry, India mock, Tall nettle, Tall vervain, Tansy ragwort, Tansy mustard, Tanweed, Thistle, Trailing crownvetch, Tumble mustard, Tumble pigweed, Velvetleaf, Venice mallow, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild mustard, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Woodsorrel, Woolly croton, Woolly morningglory, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower, pepperweed and other broadleaf weeds

SPREADERS AND CALIBRATION

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

SINGLE PASS – Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 15,000 sq ft.

CALIBRATION INSTRUCTIONS: For 3.2 pounds of product per 1,000 sq ft of turfgrass: Check your spreader on 310 sq ft and adjust to apply at the rate of 1 pound of product per 310 sq ft.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owners' Manual.

NEVER APPLY BY HAND.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

SPREADER SETTINGS:

AJUSTES DEL ESPARCIDOR:

ROTARY SPREADERS *ESPARCIDORES ROTATIVOS*	
EARTHWAY 2050	14 1/4
SCOTTS SPEEDYGREEN 1000	5 3/4
SCOTTS SPEEDYGREEN 2000	6 3/4
SCOTTS SPEEDYGREEN 3000	4 1/2
VIGORO 4000	4 3/4
VIGORO 4300	4 3/4
DROP SPREADERS *ESPARCIDORES POR GOTEO*	
SCOTTS ACCUGREEN 1000	8 1/2
SCOTTS ACCUGREEN 2000	10
SCOTTS ACCUGREEN 3000	7 1/2
VIGORO 4000	1 1/4



STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.
- If overseeding, wait 4 weeks after grass seeds sprout before applying this product.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-561-1333 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water.
Do not contaminate water when disposing of equipment washwater.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	28.0%
1.17% Ammoniacal Nitrogen	
26.83 % Urea Nitrogen*	
Available Phosphate (P_2O_5)	3.0%
Soluble Potash (K_2O)	3.0%

Derived from: Polymer-coated Urea, Polymer-coated Sulfer-coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

*Contains 6.0% slowly available nitrogen from coated urea.

F1074

Distributed and guaranteed by Spectrum Group
Div. of United Industries Corp., PO Box 142642
St. Louis, MO 63114-0642 11-7465 ©2005 UIC

LIMITED WARRANTY AND DISCLAIMER.
The distributor warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.
THE DISTRIBUTOR NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the distributor or the seller. Any damages arising from a breach of the distributor's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.
The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of the Limited Warranty and Disclaimer in any manner.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.

Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.regulatory-info-sb.com**

Ultra Turf™ is a trademark of Martin Resources, Inc. and is used under license. **CTR® is a registered trademark of United Industries Corporation, and is the name for the coated urea in this product.

EPA Reg. No. 2217-819-8845
EPA Est. Nos. 9198-OH-1 (M), 9198-OH-2 (B), 9198-AL-1 (A), 49991-TX-1 (AZ), 49991-TX-2 (BC), 45065-MN-1 (BD), 59144-GA-1 (BE), 6665-NY-1 (BF), 35512-FL-2 (BG), 8378-IN-1 (AS), 43854-MO-1 (BH), 49710-TX-1 (AT), 49710-TX-3 (AU), 3234-UT-1 (AV), 3234-AZ-1 (AW), 8660-AL-1 (BN), 8660-OH-1 (BO), 74764-CA-1 (BK), 74764-FL-1 (BL), 9404-FL-1 (AX), 35497-OR-1 (AY) Circled letter is first letter of lot number.

Visit Us On The Internet
or Call Our Customer Satisfaction Hotline

Visitenos En El Internet
Para Información En Español

vigoro.com or 1-800-561-1333

Español

ALTO PRIMERO LEA TODA LA ETIQUETA. OBSERVE TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES CUIDADOSAMENTE.

GUÍA DEL PROYECTO

Para obtener un césped más tupido, más verde y más hermoso, siga el ***Programa de Cuidados de Césped Anual de Vigoro***. ¡Es fácil de seguir y puede comenzar en cualquier momento simplemente aplicando el producto que corresponde a la temporada actual!

PRIMAVERA ▸ ▸ ▸ ▸ ▸ OTOÑO ▸ ▸ ▸

Principios de la Primavera	*Fines de la Primavera*	*Verano*	*Principios del Otoño*	*Fines del Otoño*
INHIBIDOR DE HIERBAS SILVESTRES	HERBICIDA Y FERTILIZANTE	CONTROL DE INSECTOS DEL CÉSPED	CONTROL DE INSECTOS DEL CÉSPED	WINTERIZER

Vigoro ofrece programas anuales completos de cuidado del césped para todas las áreas. Consulte a un asociado de su tienda local o visite ***Vigoro.com*** para obtener más información acerca de algún programa personalizado que sea adecuado para su área y tipo de césped.

INSTRUCCIONES DE USO

El uso de este producto de forma diferente a lo indicado en su etiqueta constituye una violación de la ley Federal. No permita que personas (que no sean el aplicador) ni mascotas ingresen a la superficie tratada durante la aplicación. No ingrese a las superficies tratadas hasta que el polvo se haya asentado. No lo utilice en césped que se cultive para la venta u otro uso comercial, ni para la producción de semilla comercial, ni con fines de investigación.

DÓNDE UTILIZARLO:

El Herbicida y Fertilizante 28-3-3 Ultra Turf™ de Vigoro® controla una amplia variedad de malezas de hoja ancha en el césped de su casa. Además, el fertilizante ofrece alimento para plantas.

Utilícelo en:

Césped Bahía, Césped Bermuda, Césped Búfalo, Ciempiés, Festuca, Poa de los prados, Ballico perenne, Césped San Agustín, Césped Zoysia

CUÁNTO APLICAR:

Para Poa de los prados, festucas, ballico perenne, césped Bermuda, césped Bahía, césped Búfalo, césped Zoysia, césped San Agustín y ciempiés:

Aplique 3.2 libras del producto por cada 1,000 pies² de hierba de césped. Es probable que sea necesaria una segunda aplicación si hay una infestación densa de malezas o ante condiciones climáticas adversas. La segunda aplicación (de 3.2 libras de producto por cada 1,000 pies²) deberá realizarse 30 días después de la primera aplicación. **No aplique este producto al césped 'Floratam' San Agustín en Florida.**

La aplicación máxima sobre el césped es de 1.1 libra de equivalente ácido 2,4-D por acre por aplicación por sitio. La cantidad máxima de aplicaciones al voleo por sitio de tratamiento es 2 por año.

Es esencial una distribución uniforme del producto sobre la superficie tratada. Escoja un día calmo para asegurar una cobertura uniforme y evitar esparcir gránulos sobre las flores, vegetales y arbustos decorativos. No se exceda en la cantidad que aplique sobre el sitio y tenga cuidado al aplicar este producto en el sistema de regado por goteo de árboles, arbustos y otras plantas decorativas.

El césped generalmente utiliza entre 1 y 4 libras de nitrógeno por cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica como se indica, este producto proporciona 0.89 libras de nitrógeno por cada 1,000 pies². Comuníquese con el Servicio de Extensión local si tiene preguntas relacionadas con un programa de abono adecuado para su región geográfica.

INSTRUCCIONES DE APLICACIÓN:

1. Corte el césped a una altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped por lo menos 1 día antes de la aplicación para mantener la humedad hasta el próximo riego (vea el paso 4).
3. Aplique sobre malezas jóvenes y que estén creciendo activamente, preferentemente por la mañana cuando hay rocío sobre el césped. Si el césped no está húmedo en el momento de la aplicación, riéguelo un poco con agua para permitir que los gránulos se adhieran y permanezcan sobre la superficie de las hojas de las malezas.
4. Después de la aplicación, no riegue el césped por 1 a 2 días. Riegue, rocíe o humedezca bien el césped 2 a 3 días después de la aplicación. No aplique si se espera lluvia dentro de los 2 días.

Malezas Controladas

Annual yellow sweetclover, Aster, Austrian fieldcress, Bedstraw, Beggarticks, Betony, Florida, Bindweed, field, Bird vetch, Bitter wintercress, Bittercress, hairy, Bitterweed, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Bloodflower milkweed, Blue lettuce, Blue vervain, Bracted plantain, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull thistle, Bullnettle, Burclover, Burdock, Burning nettle, Burweed, Buttercup, Buttonweed, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catnip, Catsear, Chickweed, common, Chickweed, mouseear, Chicory, Cinquefoil, Clover, crimson, Clover, hop, Clover, red, Clover, strawberry, Clover, sweet, Clover, white, Cockle, Cocklebur, Common mullein, Creeping jenny, Cudweed, Curly dock, Daisy, English, Daisy fleabane, Daisy, oxeye, Dandelion, Dichondra, Dogfennel, Dollarweed, False dandelion, Falseflax, False sunflower, Fiddleneck, Florida pusley, Frenchweed, Galinsoga, Goldenrod, Ground ivy, Gumweed, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Heath aster, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk vetch, Morningglory, Mouseear hawkweed, Mugwort, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oriental cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prairie sunflower, Prickly lettuce, Prickly sida, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purslane, common, Ragweed, Red sorrel, Redroot pigweed, Redstem filaree, Rough cinquefoil, Rough fleabane, Russian pigweed, Russian thistle, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smooth dock, Smooth pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny amaranth, Spiny cocklebur, Spiny sowthistle, Spotted catsear, Spotted spurge, Spurweed, Stinging nettle, Strawberry, India mock, Tall nettle, Tall vervain, Tansy ragwort, Tansy mustard, Tanweed, Thistle, Trailing crownvetch, Tumble mustard, Tumble pigweed, Velvetleaf, Venice mallow, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild mustard, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Woodsorrel, Woolly croton, Woolly morningglory, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower, pepperweed and other broadleaf weeds

ESPARCIDORES Y CALIBRACIÓN

Los esparcidores por goteo y los giratorios deberán calibrarse para permitir una aplicación uniforme del producto.

UNA SOLA PASADA – Haga funcionar el equipo en intervalos según el ancho o la franja del esparcidor. Para esparcidores por goteo, superponga los rieles de las ruedas para asegurar una cobertura completa. Cierre las aperturas del esparcidor cuando lo gire para cambiarlo de dirección o cuando lo detenga.

Cobertura: Para determinar cuánto necesita aplicar, multiplique el largo de la extensión de césped por su ancho. Esto equivale a los pies cuadrados totales. Reste las áreas sin césped (casas, jardín, etc.). Una bolsa rinde 15,000 pies².

INSTRUCCIONES DE CALIBRACIÓN: Para 3.2 libras de producto por cada 1,000 pies² de hierba de césped: Verifique que su esparcidor indique 310 pies² y ajuste para aplicar en una proporción de 1 libra por producto por cada 310 pies².

IMPORTANTE: Es importante seguir las instrucciones indicadas en este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede dañar su césped. Los ajustes del esparcidor son aproximados. Las diferencias pueden darse debido a la condición del esparcidor, la velocidad a la que se aplica el producto y la rugosidad de la superficie tratada. Asegúrese de verificar el esparcidor antes de comenzar y ajústelo si es necesario. Consulte el manual del propietario.

NUNCA APLIQUE CON LA MANO.

Para evitar que queden huecos o excesos de aplicación, comience desde el borde del terreno y luego continúe hacia el centro con pasadas paralelas uniformes. Camine a un ritmo estable de modo que aplique el producto uniformemente sobre el césped.

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO: Conserve en el envase original y guarde las bolsas que contengan el producto que quedó sin usar en un lugar bajo llave al que no tengan acceso los niños ni los animales domésticos.

ELIMINACIÓN DEL CONTENEDOR:

Si está vacío: No vuelva a utilizar este contenedor. Colóquelo en un basurero u ofrézcalo para reciclaje si el servicio está disponible.

Si está parcialmente lleno: Llame a la agencia de desechos sólidos de su zona o llame al número gratuito (1-800-CLEANUP) para obtener instrucciones acerca de cómo eliminar dichos desechos. Nunca coloque el producto que quedó sin usar en un drenaje interior o exterior.

PRECAUCIONES DE USO:

- No aplique este producto a carpetgrass, dichondra, plantas decorativas ni a césped donde existan tréboles que desee conservar.
- Escoja un día calmo para asegurar una cobertura uniforme y evitar esparcir gránulos sobre las flores, vegetales y arbustos decorativos.
- No se exceda en la cantidad que aplica sobre el césped. Cuando aplique el producto debajo de un árbol, discontinúe la aplicación y siga en línea recta al otro lado del árbol, ya que si circulara alrededor pasaría dos veces por algunas áreas y en consecuencia aplicaría demasiado producto.
- Si ha resembrado sobre césped viejo, aplique este producto 4 semanas después del brote de las semillas.

DECLARACIONES DE PRECAUCIÓN

Peligroso para Humanos y Animales Domésticos

PRECAUCIÓN. Dañino si se ingiere o se absorbe a través de la piel. Causa irritación moderada en los ojos. Evite el contacto con piel, ojos o ropa. Evite respirar el polvo. Cuando utilice este producto, use camisa de mangas largas, pantalones largos, medias y zapatos.

Luego de utilizarlo, quítese la ropa y lávela por separado antes de volver a usarla y de inmediato lávese bien con agua y jabón las manos y la piel expuesta.

Primeros Auxilios

Si penetra en los ojos:	• Mantenga los ojos abiertos y enjuague lenta y suavemente con agua durante 15 a 20 minutos. Quítese los lentes de contacto (si es que usa) después de los primeros 5 minutos, luego siga enjuagando los ojos. • Llame a un centro de toxicología o a un médico para que le prescriba un tratamiento.
Si entra en contacto con la piel o la ropa:	• Quítese la ropa contaminada. • Enjuague la piel de inmediato con abundante agua durante 15 a 20 minutos. • Llame a un centro de toxicología o a un médico para que le prescriba un tratamiento.
Si lo ingiere:	• Llame inmediatamente a un centro de toxicología o a un médico para que le prescriba un tratamiento. • Si puede tragar, beba un vaso de agua. • No provoque el vómito salvo que así se lo indiquen el centro de toxicología o el médico. • No intente suministrar algo por boca a una persona inconsciente.
Tenga a mano el contenedor o la etiqueta del producto cuando llame al centro de toxicología o al médico, o cuando busque tratamiento. También puede llamar al 1-800-561-1333 para obtener información sobre el tratamiento médico de emergencia.	

PELIGROS AMBIENTALES:

La deriva o el desplazamiento pueden afectar en forma adversa a otras plantas. No lo aplique directamente en el agua. No contamine el agua cuando elimine el agua usada para la limpieza de los equipos.

0 13013 22344 1

Vigoro
Ultra Turf™

Weed & Feed

28-3-3

Herbicida y Fertilizante

TREATS 15,000 SQ FT

TRATA 15,000 PIES²

FPO UPC
0 13013 22344 1

Vigoro
Ultra Turf™
Weed & Feed
28-3-3
Herbicida y Fertilizante

Kills Over 200 Weed Types!

¡Elimina más de 200 Tipos de Malezas!

- ***Controls Dandelion, White Clover, Plantain, Chickweed, Spurge*** *& Many Other Listed Weeds*
  Controla el Diente de León, el Trébol Blanco, el Llantén, la Pamplina, la Lechetrezna *y muchas otras Malezas Enumeradas*
- ***Greens in 72 Hours***
  Proporciona Color Verde en 72 Horas
- ***Contains Composite Timed-Release (CTR®)*****
  Nitrogen for Extended Feeding
  Contiene Fertilizante de Liberación Programada Compuesta (CTR®)**
  para una Nutrición de Mayor Duración

ACTIVE INGREDIENTS:	
2,4-D	0.64%
Mecoprop-p	0.16%
Dicamba	0.03%
OTHER INGREDIENTS:	99.17%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
12.80 lbs. 2, 4-dichlorophenoxyacetic acid equivalent per ton.
3.20 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
0.60 lbs. dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.
Isomer Specific by AOAC Methods
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

CAS Registry Numbers:
2,4-D CAS# 94-75-7
Mecoprop-p CAS# 16484-77-8
Dicamba CAS# 1918-00-9

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements and First Aid.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN Consulte el panel posterior para obtener Declaraciones de Precaución y Primeros Auxilios adicionales.

Net Wt 48 lb (21.7 kg)

TREATS 15,000 SQ FT
Trata 15,000 pies²

APPLY ANYTIME WEEDS ARE ACTIVELY GROWING
Aplique en cualquier momento en que las malezas estén creciendo activamente

Vigoro Ultra Turf™

Weed & Feed
28-3-3
Herbicida y Fertilizante

Weed & Feed 28-3-3 Herbicida y Fertilizante
TREATS 15,000 SQ FT TRATA 15,000 PIES²

FPO UPC
0 13013 22344 1

English

STOP **READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

PROJECT GUIDE

For a thicker, greener, more beautiful lawn, follow ***Vigoro's Annual Lawn Care Program***. It's easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!

SPRING ▸ ▸ ▸ ▸ ▸ **FALL** ▸ ▸ ▸

Early Spring	*Late Spring*	*Summer*	*Early Fall*	*Late Fall*
CRABGRASS PREVENTER	WEED & FEED	LAWN INSECT CONTROL	LAWN FERTILIZER	WINTERIZER

Vigoro offers complete annual lawn care programs for all areas. See your local store associate or visit ***Vigoro.com*** for more information about a customized program that is right for your area and turf type.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

Vigoro® Ultra Turf™ Weed & Feed 28-3-3 controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

For Use On:

Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, Fescue, Kentucky bluegrass, Perennial ryegrass, St. Augustinegrass, Zoysiagrass

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass:

Apply 3.2 lb of product per 1,000 sq ft of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 lb of product per 1,000 sq ft should be used 30 days after the first application. **Do not apply this product to 'Floratam' St. Augustinegrass in Florida.**

The maximum application rate to turf is 1.1 lb of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

A lawn typically utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.89 lb of nitrogen per 1,000 sq ft. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application. Do not apply if rain is expected within 2 days.

Weeds Controlled

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Clover, sweet
Clover, white
Cockle
Cocklebur
Common mullein
Creeping jenny
Cudweed
Curly dock
Daisy, English
Daisy fleabane
Daisy, oxeye
Dandelion
Dichondra
Dogfennel
Dollarweed
False dandelion
Falseflax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed and other broadleaf weeds

SPREADER SETTINGS:

Español

ALTO **PRIMERO LEA TODA LA ETIQUETA. OBSERVE TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES CUIDADOSAMENTE.**

GUÍA DEL PROYECTO

Para obtener un césped más tupido, más verde y más hermoso, siga el ***Programa de Cuidados de Césped Anual de Vigoro***. ¡Es fácil de seguir y puede comenzar en cualquier momento simplemente aplicando el producto que corresponde a la temporada actual!

PRIMAVERA ▸ ▸ ▸ ▸ ▸ **OTOÑO** ▸ ▸ ▸

Principios de la Primavera	*Fines de la Primavera*	*Verano*	*Principios del Otoño*	*Fines del Otoño*
INHIBIDOR DE HIERBAS SILVESTRES	HERBICIDA Y FERTILIZANTE	CONTROL DE INSECTOS DEL CÉSPED	CONTROL DE INSECTOS DEL CÉSPED	WINTERIZER

Vigoro ofrece programas anuales completos de cuidado del césped para todas las áreas. Consulte a un asociado de su tienda local o visite ***Vigoro.com*** para obtener más información acerca de algún programa personalizado que sea adecuado para su área y tipo de césped.

INSTRUCCIONES DE USO

El uso de este producto de forma diferente a lo indicado en su etiqueta constituye una violación de la ley Federal. No permita que personas (que no sean el aplicador) ni mascotas ingresen a la superficie tratada durante la aplicación. No ingrese a las superficies tratadas hasta que el polvo se haya asentado. No lo utilice en césped que se cultive para la venta u otro uso comercial, ni para la producción de semilla comercial, ni con fines de investigación.

DÓNDE UTILIZARLO:

El Herbicida y Fertilizante 28-3-3 Ultra Turf™ de Vigoro® controla una amplia variedad de malezas de hoja ancha en el césped de su casa. Además, el fertilizante ofrece alimento para plantas.

Utilícelo en:

Césped Bahía, Césped Bermuda, Césped Búfalo, Ciempiés, Festuca, Poa de los prados, Ballico perenne, Césped San Agustín, Césped Zoysia

CUÁNTO APLICAR:

Para Poa de los prados, festucas, ballico perenne, césped Bermuda, césped Bahía, césped Búfalo, césped Zoysia, césped San Agustín y ciempiés:

Aplique 3.2 libras del producto por cada 1,000 pies² de hierba de césped. Es probable que sea necesaria una segunda aplicación si hay una infestación densa de malezas o ante condiciones climáticas adversas. La segunda aplicación (de 3.2 libras de producto por cada 1,000 pies²) deberá realizarse 30 días después de la primera aplicación. **No aplique este producto al césped 'Floratam' San Agustín en Florida.**

La aplicación máxima sobre el césped es de 1.1 libra de equivalente ácido 2,4-D por acre por aplicación por sitio. La cantidad máxima de aplicaciones al voleo por sitio de tratamiento es 2 por año.

Es esencial una distribución uniforme del producto sobre la superficie tratada. Escoja un día calmo para asegurar una cobertura uniforme y evitar esparcir gránulos sobre las flores, vegetales y arbustos decorativos. No se exceda en la cantidad que aplique sobre el sitio y tenga cuidado al aplicar este producto en el sistema de regado por goteo de árboles, arbustos y otras plantas decorativas.

El césped generalmente utiliza entre 1 y 4 libras de nitrógeno por cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica como se indica, este producto proporciona 0.89 libras de nitrógeno por cada 1,000 pies². Comuníquese con el Servicio de Extensión local si tiene preguntas relacionadas con un programa de abono adecuado para su región geográfica.

INSTRUCCIONES DE APLICACIÓN:

1. Corte el césped a una altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped por lo menos 1 día antes de la aplicación para mantener la humedad hasta el próximo riego (vea el paso 4).
3. Aplique sobre malezas jóvenes y que estén creciendo activamente, preferentemente por la mañana cuando hay rocío sobre el césped. Si el césped no está húmedo en el momento de la aplicación, riéguelo un poco con agua para permitir que los gránulos se adhieran y permanezcan sobre la superficie de las hojas de las malezas.
4. Después de la aplicación, no riegue el césped por 1 a 2 días. Riegue, rocíe o humedezca bien el césped 2 a 3 días después de la aplicación. No aplique si se espera lluvia dentro de los 2 días.

Malezas Controladas

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Clover, sweet
Clover, white
Cockle
Cocklebur
Common mullein
Creeping jenny
Cudweed
Curly dock
Daisy, English
Daisy fleabane
Daisy, oxeye
Dandelion
Dichondra
Dogfennel
Dollarweed
False dandelion
Falseflax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed and other broadleaf weeds

Bull thistle Cudweed Hoary cress Pennycress Slender plantain Western clematis

SPREADERS AND CALIBRATION

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

SINGLE PASS — Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 15,000 sq ft.

CALIBRATION INSTRUCTIONS: For 3.2 pounds of product per 1,000 sq ft of turfgrass: Check your spreader on 310 sq ft and adjust to apply at the rate of 1 pound of product per 310 sq ft.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owners' Manual.

NEVER APPLY BY HAND.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

SPREADER SETTINGS:

AJUSTES DEL ESPARCIDOR:

ROTARY SPREADERS ***ESPARCIDORES ROTATIVOS***	
EARTHWAY 2050	14 1/4
SCOTTS SPEEDYGREEN 1000	5 3/4
SCOTTS SPEEDYGREEN 2000	6 3/4
SCOTTS SPEEDYGREEN 3000	4 1/2
VIGORO 4000	4 3/4
VIGORO 4300	4 3/4
DROP SPREADERS ***ESPARCIDORES POR GOTEO***	
SCOTTS ACCUGREEN 1000	8 1/2
SCOTTS ACCUGREEN 2000	10
SCOTTS ACCUGREEN 3000	7 1/2
VIGORO 4000	1 1/4

Bull thistle Cudweed Hoary cress Pennycress Slender plantain Western clematis

ESPARCIDORES Y CALIBRACIÓN

Los esparcidores por goteo y los giratorios deberán calibrarse para permitir una aplicación uniforme del producto.

UNA SOLA PASADA — Haga funcionar el equipo en intervalos según el ancho o la franja del esparcidor. Para esparcidores por goteo, superponga los rieles de las ruedas para asegurar una cobertura completa. Cierre las aperturas del esparcidor cuando lo gire para cambiarlo de dirección o cuando lo detenga.

Cobertura: Para determinar cuánto necesita aplicar, multiplique el largo de la extensión de césped por su ancho. Esto equivale a los pies cuadrados totales. Reste las áreas sin césped (casas, jardín, etc.). Una bolsa rinde 15,000 pies^2.

INSTRUCCIONES DE CALIBRACIÓN: Para 3.2 libras de producto por cada 1,000 pies^2 de hierba de césped: Verifique que su esparcidor indique 310 pies^2 y ajuste para aplicar en una proporción de 1 libra por producto por cada 310 pies^2.

IMPORTANTE: Es importante seguir las instrucciones indicadas en este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede dañar su césped. Los ajustes del esparcidor son aproximados. Las diferencias pueden darse debido a la condición del esparcidor, la velocidad a la que se aplica el producto y la rugosidad de la superficie tratada. Asegúrese de verificar el esparcidor antes de comenzar y ajústelo si es necesario. Consulte el manual del propietario.

NUNCA APLIQUE CON LA MANO.

Para evitar que queden huecos o excesos de aplicación, comience desde el borde del terreno y luego continúe hacia el centro con pasadas paralelas uniformes. Camine a un ritmo estable de modo que aplique el producto uniformemente sobre el césped.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.
- If overseeding, wait 4 weeks after grass seeds sprout before applying this product.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-561-1333 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water.
Do not contaminate water when disposing of equipment washwater.

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO: Conserve en el envase original y guarde las bolsas que contengan el producto que quedó sin usar en un lugar bajo llave al que no tengan acceso los niños ni los animales domésticos.

ELIMINACIÓN DEL CONTENEDOR:

Si está vacío: No vuelva a utilizar este contenedor. Colóquelo en un basurero u ofrézcalo para reciclaje si el servicio está disponible.

Si está parcialmente lleno: Llame a la agencia de desechos sólidos de su zona o llame al número gratuito (1-800-CLEANUP) para obtener instrucciones acerca de cómo eliminar dichos desechos. Nunca coloque el producto que quedó sin usar en un drenaje interior o exterior.

PRECAUCIONES DE USO:

- No aplique este producto a carpetgrass, dichondra, plantas decorativas ni a césped donde existan tréboles que desee conservar.
- Escoja un día calmo para asegurar una cobertura uniforme y evitar esparcir gránulos sobre las flores, vegetales y arbustos decorativos.
- No se exceda en la cantidad que aplica sobre el césped. Cuando aplique el producto debajo de un árbol, discontinúe la aplicación y siga en línea recta al otro lado del árbol, ya que si circulara alrededor pasaría dos veces por algunas áreas y en consecuencia aplicaría demasiado producto.
- Si ha resembrado sobre césped viejo, aplique este producto 4 semanas después del brote de las semillas.

DECLARACIONES DE PRECAUCIÓN

Peligroso para Humanos y Animales Domésticos

PRECAUCIÓN. Dañino si se ingiere o se absorbe a través de la piel. Causa irritación moderada en los ojos. Evite el contacto con piel, ojos o ropa. Evite respirar el polvo. Cuando utilice este producto, use camisa de mangas largas, pantalones largos, medias y zapatos.

Luego de utilizarlo, quítese la ropa y lávela por separado antes de volver a usarla y de inmediato lávese bien con agua y jabón las manos y la piel expuesta.

Primeros Auxilios

Si penetra en los ojos:	• Mantenga los ojos abiertos y enjuague lenta y suavemente con agua durante 15 a 20 minutos. Quítese los lentes de contacto (si es que usa) después de los primeros 5 minutos, luego siga enjuagando los ojos. • Llame a un centro de toxicología o a un médico para que le prescriba un tratamiento.
Si entra en contacto con la piel o la ropa:	• Quítese la ropa contaminada. • Enjuague la piel de inmediato con abundante agua durante 15 a 20 minutos. • Llame a un centro de toxicología o a un médico para que le prescriba un tratamiento.
Si lo ingiere:	• Llame inmediatamente a un centro de toxicología o a un médico para que le prescriba un tratamiento. • Si puede tragar, beba un vaso de agua. • No provoque el vómito salvo que así se lo indiquen el centro de toxicología o el médico. • No intente suministrar algo por boca a una persona inconsciente.
Tenga a mano el contenedor o la etiqueta del producto cuando llame al centro de toxicología o al médico, o cuando busque tratamiento. También puede llamar al 1-800-561-1333 para obtener información sobre el tratamiento médico de emergencia.	

PELIGROS AMBIENTALES:

La deriva o el desplazamiento pueden afectar en forma adversa a otras plantas. No lo aplique directamente en el agua. No contamine el agua cuando elimine el agua usada para la limpieza de los equipos.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	28.0%
1.17% Ammoniacal Nitrogen	
26.83 % Urea Nitrogen*	
Available Phosphate (P_2O_5)	3.0%
Soluble Potash (K_2O)	3.0%

Derived from: Polymer-coated Urea, Polymer-coated Sulfer-coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

*Contains 6.0% slowly available nitrogen from coated urea.

F1074

Distributed and guaranteed by Spectrum Group
Div. of United Industries Corp., PO Box 142642
St. Louis, MO 63114-0642 11-7465 ©2005 UIC

LIMITED WARRANTY AND DISCLAIMER.
The distributor warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.
THE DISTRIBUTOR NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the distributor or the seller. Any damages arising from a breach of the distributor's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.
The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of the Limited Warranty and Disclaimer in any manner.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.

Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.regulatory-info-sb.com**

Ultra Turf™ is a trademark of Martin Resources, Inc. and is used under license. **CTR® is a registered trademark of United Industries Corporation, and is the name for the coated urea in this product.

Visit Us On The Internet
or Call Our Customer Satisfaction Hotline

Visitenos En El Internet
Para Información En Español

vigoro.com or 1-800-561-1333

EPA Reg. No. 2217-819-8845
EPA Est. Nos. 9198-OH-1 (M), 9198-OH-2 (B), 9198-AL-1 (A), 49991-TX-1 (AZ), 49991-TX-2 (BC), 45065-MN-1 (BD), 59144-GA-1 (BE), 6665-NY-1 (BF), 35512-FL-2 (BG), 8378-IN-1 (AS), 43854-MO-1 (BH), 49710-TX-1 (AT), 49710-TX-3 (AU), 3234-UT-1 (AV), 3234-AZ-1 (AW), 8660-AL-1 (BN), 8660-OH-1 (BO), 74764-CA-1 (BK), 74764-FL-1 (BL), 9404-FL-1 (AX), 35497-OR-1 (AY) Circled letter is first letter of lot number.